

Real-analytic, volume-preserving actions of lattices on 4-manifolds

Benson Farb and Peter Shalen *

February 1, 2008

Abstract

We prove that any real-analytic, volume-preserving action of a lattice Γ in a simple Lie group with $\mathbf{Q}$ -rank(Γ) ≥ 7 on a closed 4-manifold of nonzero Euler characteristic factors through a finite group action.

1 Results

Zimmer conjectured in [Zi1] that the standard action of $\mathrm{SL}(n, \mathbf{Z})$ on the n -torus is minimal in the following sense:

Conjecture 1 (Zimmer). *Any smooth, volume-preserving action of any finite-index subgroup $\Gamma < \mathrm{SL}(n, \mathbf{Z})$ on a closed r -manifold factors through a finite group action if $n > r$.*

While Conjecture 1 has been proved for actions which also preserve an extra geometric structure such as a pseudo-Riemannian metric (see, e.g. [Zi1]), almost nothing is known in the general case. For $r = 2$ and $n > 4$, the conjecture was proved for real-analytic actions in [FS1]. Quite recently, Polterovich [Po] has brought ideas from symplectic topology to the problem, using these to give a proof of Conjecture 1 for orientable surfaces of genus > 1 . In [FS2] we will point out how his methods actually prove Conjecture 1 for the torus as well. For $r = 3$, Conjecture 1 is known only in some special cases where Γ contains some torsion and the action is real-analytic (see [FS1]).

In this note we prove the following result.

*Both authors are supported in part by the NSF.

Theorem 2 (Actions on 4-manifolds with $\chi(M) \neq 0$). *Let Γ be a lattice in a simple Lie group G such that $\mathbf{Q}\text{-rank}(\Gamma) \geq 7$. Then any real-analytic, volume-preserving action of Γ on a closed 4-manifold of nonzero Euler characteristic factors through a finite group action.*

In particular, Theorem 2 applies to any finite-index subgroup of $\mathrm{SL}(n, \mathbf{Z})$, $n > 7$.

The main ingredient in the proof of Theorem 2 is Theorem 7.1 of [FS1] on real-analytic actions which preserve a volume form. This theorem, which is the most difficult result in [FS1], gives a *codimension-two* invariant submanifold for centralizers of elements with fixed-points. One can then apply results of [FS1] and [Re], which show that real-analytic (not necessarily area-preserving) actions of certain lattices on 2-dimensional manifolds must factor through finite groups.

For the case of symplectic actions, some further progress on Conjecture 1 can be found in [Po] and [FS2].

2 Proof of Theorem 2

Before giving the proof of Theorem 2, we will need two algebraic properties of lattices with large $\mathbf{Q}$ -rank.

Proposition 3 (Some algebraic properties of lattices). *Let Γ be a lattice in a simple algebraic group over $\mathbf{Q}$. Then the following hold:*

1. *If $d = \mathbf{Q}\text{-rank}(\Gamma) \geq 7$ then Γ contains commuting subgroups A and B which are isomorphic to irreducible lattices with $\mathbf{Q}$ -ranks 2 and $d - 3$ respectively.*
2. *If $\mathbf{Q}\text{-rank}(\Gamma) \geq 4$ then Γ contains a torsion-free nilpotent subgroup which is not metabelian.*

Proof. The proof of the first statement is similar to that of Proposition 2.1 of [FS1]. By Margulis's Arithmeticity Theorem (see, e.g., [Zi2], Theorem 6.1.2), Γ is commensurate with the group of $\mathbf{Z}$ -points of a simple algebraic group G defined over $\mathbf{Q}$. Hence without loss of generality we can assume that Γ itself is the group of $\mathbf{Z}$ -points in such a group G .

Since G is simple, the root system Φ of G is irreducible, and the Dynkin diagram determined Φ therefore appears in the list given in Section 11.4 of [Hu]. By going through this list, one sees that in every case where $d \geq 7$, one may “erase a vertex v ” of the diagram to obtain a graph with 2 components:

one with two vertices and another which is a Dynkin diagram with at least $d - 3$ vertices. Let G_1 and G_2 be the root subgroups corresponding to these two components of the Dynkin diagram. Then the group of $\mathbf{Q}$ -points of G_1 has $\mathbf{Q}$ -rank at least 2, the group of $\mathbf{Q}$ -points of G_2 has $\mathbf{Q}$ -rank at least $d - 3$, and G_1 commutes with G_2 .

Now $\Gamma_i = \Gamma \cap G_i$ is an arithmetic lattice in G_i for $i = 1, 2$, since by a theorem of Borel-Harish-Chandra (see, e.g. [Zi2]) the $\mathbf{Z}$ -points of an algebraic group defined over $\mathbf{Q}$ form a lattice in the group of $\mathbf{R}$ -points. Then $A = \Gamma_1$ and $B = \Gamma_2$ have the required properties.

To prove the second statement, note that since $\mathbf{Q}\text{-rank}(\Gamma) \geq 4$, we can find a connected, nilpotent Lie subgroup N which is defined over $\mathbf{Q}$ and has derived length 3, i.e. is not metabelian. As $\Gamma \cap N$ is the group of $\mathbf{Z}$ -points of the $\mathbf{Q}$ -group N , it is a lattice in N , and in particular is Zariski-dense in N . Hence $\Gamma \cap N$ is nilpotent and has no metabelian subgroup of finite index. As $\Gamma \cap N$ must have a torsion-free subgroup of finite index, the assertion follows. $\diamond$

We now turn to the proof of Theorem 2. Let M be a closed 4-manifold with nonzero Euler characteristic. Let Γ be an irreducible lattice in a simple Lie group G , and assume $d = \mathbf{Q}\text{-rank}(\Gamma) \geq 7$. By part (1) of Proposition 3, Γ contains commuting subgroups A and B which are isomorphic to irreducible lattices with $\mathbf{Q}$ -ranks 2 and $d - 3 \geq 4$ respectively.

Let γ_0 be any infinite order element of A . By an old theorem of Fuller [Fu], any homeomorphism of a closed manifold of nonzero Euler characteristic has a periodic point; the proof is an application of the Lefschetz fixed-point theorem and basic number theory. Hence some positive power γ of γ_0 has a fixed point.

We will also need the following two facts. First, since $\mathbf{Q}\text{-rank}(B) \geq d - 3 \geq 4$, it follows from Margulis's Superrigidity Theorem that any representation of B into $\text{GL}(4, \mathbf{R})$ has finite image. Second, since Γ is a lattice in a simple Lie group G with $\mathbf{R}\text{-rank}(G) \geq 2$, the Margulis Finiteness Theorem (see, e.g., Theorem 8.1 of [Zi2]) gives that Γ is *almost simple* in the sense that any normal subgroup of Γ must be finite or of finite index.

We are now in a position to apply Theorem 7.1 of [FS1]. For the reader's convenience we recall the statement here. We say that a group action $\rho : \Gamma \rightarrow \text{Diff}(M)$ is *infinite* if ρ has infinite image.

Theorem 7.1 of [FS1]: *Let Γ be an almost simple group. Suppose we are given an infinite, volume-preserving, real-analytic action of Γ on a closed,*

connected n -manifold M . Suppose further that Γ contains commuting subgroups A and B with the following properties:

- There exists an element $\gamma \in A$, noncentral in Γ , having a fixed point in M .
- A is isomorphic to an irreducible lattice of $\mathbf{Q}$ -rank ≥ 2 .
- B is noncentral in Γ .
- Any representation of any finite-index subgroup of B in $\mathrm{GL}(n, \mathbf{R})$ has finite image.

Then there is a nonempty, connected, real-analytic submanifold $W \subset M$ of codimension at least 2 which is invariant under a finite-index subgroup B' of B . Furthermore, the action of this subgroup on W is infinite.

Remark. The action of B' on the surface W produced by this theorem is NOT necessarily area preserving.

We now conclude the proof of Theorem 2. Since B' is a lattice in a simple Lie group and $\mathbf{Q}\text{-rank}(B') \geq 4$, it follows from part (2) of Proposition 3 that B' contains a torsion-free nilpotent subgroup H which is not metabelian. But Rebelo [Re] showed that any nilpotent group of real-analytic diffeomorphisms of a compact, oriented surface must be metabelian. It follows that the action of H on W is not effective.

Since H is torsion-free, there is an infinite-order element of $H \leq B'$ which acts trivially on W . Since B' has finite index in the almost simple group B , and hence is almost simple, some finite index subgroup C of B' acts trivially on W ; in particular C has a global fixed point in M . Since C is isomorphic to a lattice of $\mathbf{Q}$ -rank at least 4, by Lemma 3.2 of [FS1] we have that a finite index subgroup D of C acts trivially on M . Since Γ is almost simple, it follows that some finite index normal subgroup of Γ acts trivially on M , and we are done. $\diamond$

References

- [FS1] B. Farb and P. Shalen, Real-analytic actions of lattices, *Inventiones Math.*, Vol. 135 (1998), no. 2, p.273-296.
- [FS2] B. Farb and P. Shalen, Lattice actions on symplectic 4-manifolds, in preparation.

- [Fu] F.B. Fuller, The existence of periodic points, *Annals of Math.*, Vol. 57, No. 2 (1953), p.229-230.
- [Hu] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*, GTM 9, third printing, Springer-Verlag, 1972.
- [Po] L. Polterovich, Growth of maps, distortion in groups and symplectic geometry, December 2000 preprint.
- [Re] J. Rebelo, On nilpotent groups of real analytic diffeomorphisms of the torus, *C.R. Acad. Sci. Paris*, t. 331, No. 1 (2000), p.317-322.
- [Zi1] R. Zimmer, Actions of semisimple groups and discrete subgroups, *Proc. I.C.M.*, Berkeley 1986, p.1247-1258.
- [Zi2] R. Zimmer, *Ergodic Theory and Semisimple Groups*, Monographs in Math., Vol. 81, Birkhäuser, 1984.

Benson Farb:

Dept. of Mathematics, University of Chicago

5734 University Ave.

Chicago, IL 60637

E-mail: farb@math.uchicago.edu

Peter Shalen:

Dept. of Mathematics, University of Illinois at Chicago

Chicago, IL 60680

E-mail: shalen@math.uic.edu